

Scheme for preparation of nonmaximal entanglement between two atomic ensembles

Peng Xue*, and Guang-Can Guo†

Key Laboratory of Quantum Information, University of Science and Technology of China, Hefei 230026, P. R. China

We propose an experimentally feasible scheme to generate nonmaximal entanglement between two atomic ensembles. The degree of entanglement is readily tunable. The scheme involves laser manipulation of atomic ensembles, adjustable quarter- and half-wave plates, beam splitter, polarizing beam splitters, and single-photon detectors, and well fits the status of the current experimental technology. Finally we use the nonmaximally entangled state of ensembles to demonstrate quantum nonlocality by detecting the Clauser-Horne-Shimony-Holt inequality.

PACS number(s): 03.65.Ud, 03.67.-a, 42.50.Gy, 42.50.-p

Quantum entanglement is one of the most striking features of quantum mechanics. The recent surge of interest and progress in quantum information theory allows one to take a more positive view of entanglement and regard it as an essential resource for many ingenious applications such as quantum computation [1–3], quantum teleportation [4,5], superdense coding [6], and quantum cryptography [7–10]. The technology of generation and manipulation of bipartite or multipartite entangled states has been realized in some systems [11–18]. In most of the above schemes, the subsystems are taken as single-particle. Remarkably, Lukin and Duan *et al.* have proposed some schemes [19–22] for preparation of entanglement, in which atomic ensembles with a large number of identical atoms are used as the basic system. For example, one can use atomic ensembles for generation of substantial spin squeezing [23] and continuous variable entanglement [20,24], and for efficient preparation of Einstein-Podolsky-Rosen (EPR) [21], Greenberger-Horne-Zeilinger (GHZ) type of maximally entangled states [22] and W class of maximally entangled states [25]. The schemes have some special advantages compared with other quantum information schemes based on the control of single particles [26].

In all experimental efforts, it is hard to vary the degree of entanglement, and to produce nonmaximally entangled states without compromising the purity of the state. Nonmaximally entangled states have been shown to reduce the required detector efficiencies for loophole-free tests of Bell inequalities [27], as well as allowing logical arguments that demonstrate the nonlocality of quantum mechanics without inequalities [28]. Recently, we have

proposed some schemes for applications of nonmaximal entanglement [9,10,29]. Up to now, nonmaximally entangled states have been deterministically generated with an ion-trap [30], and with a spontaneous down converter (SPDC) [31–33].

Here, we describe an experimental scheme of preparing nonmaximally entangled states based on Raman type laser manipulation of the atomic ensembles and single photon detection which postselects the desired state in a probabilistic fashion.

The basic element of this scheme is an ensemble of many identical alkali atoms with a Raman type Λ -level configuration shown as Fig. 1, the experimental realization of which can be either a room-temperature dilute atomic gas [24,34] or a sample of cold trapped atoms [35,36]. We continue to use the symbols and corresponding definitions in Refs. [21,22]. The collective atomic operator is defined as

$$s = \left(1/\sqrt{N_a}\right) \sum_{i=1}^{N_a} |g\rangle_i \langle s|, \quad (1)$$

where $N_a \gg 1$ is the total atom number. The Raman transition $|g\rangle \rightarrow |e\rangle$ is coupled by the classical laser and the forward scattered Stokes light comes from the transition $|e\rangle \rightarrow |s\rangle$ [21]. The scheme for preparation nonmaximally entangled states works in the following way (see Fig. 2). Here we choose a Λ configuration of atomic states of ^{87}Rb by way of example, which is coupled by a pair of optical fields.

There are two light fields with the Rabi frequencies Ω and ω , respectively, which couple pairs of Zeeman sublevels of electronic ground state $5S_{1/2}$ ^{87}Rb atoms ($|g\rangle, |s\rangle$), with magnetic quantum numbers differing by two, via the excited $5P_{1/2}$ state [34,37]. In this case $|g\rangle$ and $|s\rangle$ of the simplified three-level model correspond, respectively, to $|F=2, M_F=-2\rangle$ and $|F=2, M_F=0\rangle$. The atoms in the ensembles are initially prepared to the ground state $|g\rangle$ through optical pumping. The two ensembles 1 and 2 are illuminated by the synchronized classical laser pulses of right circularly polarized (σ_-) light. The excitations $5S_{1/2} F=2, M_F=-2 \rightarrow 5P_{1/2} F=1, M_F=-1$ can be transferred to optical excitations. Assume that the light-atom interaction time t_0 is short so that the mean photon number in the forward scattered Stokes light is much smaller than 1. It is defined in Ref. [21] an effective single-mode bosonic operator a for this Stokes pulse. The whole state of the atomic collective mode and the forward scattering Stokes mode can be

*Email address: xuepeng@mail.ustc.edu.cn

†Email address: gcguo@ustc.edu.cn

written as

$$|\varphi\rangle = |vac\rangle_a |vac\rangle_p + \sqrt{p_c} s^+ a^+ |vac\rangle_a |vac\rangle_p + o(p_c), \quad (2)$$

where $|vac\rangle_a$ and $|vac\rangle_p$ denote the vacuum states of atomic ensembles and Stokes light respectively, and p_c is the small excitation probability [21]. The forward scattered Stokes pulses of left circularly polarized (σ_+) light from both ensembles are combined at a 50%–50% beam splitter (BS) and a single-photon detector click in one of the four detectors, after the quarter-and half-wave plates (QWP and HWP), and polarizing beam splitter (PBS). Adjustable QWP, HWP and PBS allow polarization analysis in any basis, i.e., at any position on the Poincare sphere [33,38].

Nonmaximally entangled states are produced simply by adjusting the relative inclination between the optic axes of QWP and HWP. After some filters which filter out the pumping laser pulses, by rotating the optic axis of QWP, the Stokes light is turned to linearly polarized photon and the orientation of the linear polarizer lies in the vertical plane. For an inclination θ_i between the optic axis of HWP and the orientation of the linear polarizer of the Stokes light in arm i ($i = 1, 2$), the Stokes photon is $\sin 2\theta_i |H\rangle + e^{i\phi} \cos 2\theta_i |V\rangle$, where H and V , respectively, represent the horizontal and vertical polarizations of the photon, and $\theta_1 = \frac{\pi}{4} - \theta_2$. Then the Stokes pulses in both arms are combined at the BS and the output light goes through a PBS, respectively, and a single-photon detector click in one of the four detectors D1, D2, D3 and D4 measures the combination radiation from the samples A^+A or A'^+A' . Here,

$$A = \alpha a_1 + e^{i\phi_{12}} \beta a_2, \quad (3)$$

or

$$A' = \beta a_1 + e^{i\phi_{12}} \alpha a_2, \quad (4)$$

where $\phi_{12} = \phi_2 - \phi_1$ is a difference of the phase shift which is fixed by the optical channel connecting the two atomic ensembles, and $\alpha = \sin 2\theta_1$, $\beta = \cos 2\theta_1$. That is, if D1 or D3 clicks, the two ensembles are entangled in the form

$$|\psi\rangle_{12} = (\alpha s_1^+ + e^{i\phi_{12}} \beta s_2^+) |vac\rangle_{12}; \quad (5)$$

if D2 or D4 clicks, the state of the ensembles is

$$|\psi'\rangle_{12} = (\beta s_1^+ + e^{i\phi_{12}} \alpha s_2^+) |vac\rangle_{12}. \quad (6)$$

The subscripts 1 and 2 are used to distinguish the atomic ensemble $E1$ and $E2$ (seeing Fig. 2). If one excitation is registered, we succeed to entangle the two ensembles in a nonmaximally entangled state. Otherwise, we need to repeat the above steps until we get a click in one of the four detectors.

Now, we consider the efficiency of this scheme, which is usually described by the total generation time. The preparation based on the Raman driving $|g\rangle \rightarrow |e\rangle \rightarrow |s\rangle$, succeeds with a controllable probability p_c for each Raman driving pulse, and needs to be repeated in average $1/p_c$ times for the final successful state generation. In the generation process, the dominant noise is the photon loss, which includes the contributions from the channel attenuation, the spontaneous emissions in the atomic ensembles, the coupling inefficiency of Stokes light into and out of the channel, and the inefficiency of the single-photon detectors which can not perfectly distinguish between one and two photons. All the above noise is described by an overall loss probability η . Due to the noise, the total generation time is represented by $T \sim t_0 / [(1 - \eta) p_c]$, where t_0 is the light-atom interaction time.

Also with the noise, the state of the ensembles is actually described by

$$\rho_{12} = \frac{1}{c+1} (c |vac\rangle_{12} \langle vac| + |\psi\rangle_{12} \langle \psi|), \quad (7)$$

and

$$\rho'_{12} = \frac{1}{c+1} (c |vac\rangle_{12} \langle vac| + |\psi'\rangle_{12} \langle \psi'|), \quad (8)$$

where the vacuum coefficient c is basically given by the conditional probability for the inherent mode-mismatching noise contribution [26].

Since the nonmaximally entangled states shown in Eqs. (5) and (6) are entangled in the Fock basis, it is experimentally hard to do certain single-bit rotation. In the following we will show how the nonmaximally entangled states can be used to realize the communication protocols, such as the CHSH detection, with simple experimental configurations.

The first step is to share an EPR type of entangled state [21]

$$|\Psi_\phi\rangle_{L_1 R_1} = (s_{L_1}^+ + e^{i\phi} s_{R_1}^+) / \sqrt{2} |vac\rangle_{L_1 R_1} \quad (9)$$

between two distant ensembles L_1 and R_1 , and the presence of the noise modifies the projected state of the ensembles to

$$\rho_{L_1 R_1} = \frac{1}{c+1} (c |vac\rangle_{L_1 R_1} \langle vac| + |\Psi_\phi\rangle_{L_1 R_1} \langle \Psi_\phi|). \quad (10)$$

The ensembles L_2 and R_2 are prepared in a nonmaximally entangled state $\rho_{L_2 R_2}$ shown in Eq. (7). The ϕ -parameters in $\rho_{L_1 R_1}$ and $\rho_{L_2 R_2}$ are the same provided that the two states are established over the same stationary channels.

A basis of the “polarization” qubit (in analogy to the language for photons) can be defined from the states $|H\rangle_i = s_{i1}^+ |vac\rangle_{i1i2}$, $|V\rangle_i = s_{i2}^+ |vac\rangle_{i1i2}$ ($i = L, R$).

Single-bit rotations in this basis can be done using the phase shift ϕ_i together with the corresponding beam splitter operation with the rotation angle $\theta_i = \phi_i/2$ similarly to the manipulations shown in Ref. [21].

The four ensembles are illuminated by the synchronized classical laser pulses with the frequency ω . If the ensemble is in the metastable state after the repumping pulse, the transition $|e\rangle \rightarrow |s\rangle$ will occur *determinately*. We register only the coincidences of the two-side detectors, so the protocol is successful only if there is a click on each side. Under this condition, the vacuum components in the entangled states and the state $s_{L_1}^+ s_{L_2}^+ |vac\rangle_{L_1 L_2 R_1 R_2}$ and $s_{R_1}^+ s_{R_2}^+ |vac\rangle_{L_1 L_2 R_1 R_2}$ have no contributions to the experimental results. Then, for the measurement scheme shown by Fig. 3, the state $\rho_{L_1 R_1} \otimes \rho_{L_2 R_2}$ is effectively equivalent to the following “polarization” nonmaximally entangled (PNE) state

$$|\psi\rangle_{PNE} = (\alpha s_{L_2}^+ s_{R_1}^+ + \beta s_{L_1}^+ s_{R_2}^+) |vac\rangle_{L_1 L_2 R_1 R_2}. \quad (11)$$

The success probability for the projection is given by $p = 1/[4(c+1)^2]$.

Now, it is clear how to do the Clauser-Horne-Shimony-Holt (CHSH) inequality detection [39]. We define the measurement results to be 1 if D1 or D3 clicks, and -1 if D2 or D4 clicks. Then the quantity

$$E(\phi_L, \phi_R) = P_{D_1 D_3} + P_{D_2 D_4} - P_{D_1 D_4} - P_{D_2 D_3} \quad (12)$$

is the corresponding coefficient of the measurements performed by Side L in the basis rotated by $\phi_L/2$ and by Side R in the basis rotated by $\phi_R/2$. According to the quantum rules

$$E(\phi_L, \phi_R) = 4\alpha^2\beta^2 \cos(\phi_L - \phi_R), \quad (13)$$

one can define the quantity S composed of the correlation coefficients for which both sides used analysis (phase shift ϕ_i) of different orientation

$$S = E(\phi_L^1, \phi_R^3) + E(\phi_L^1, \phi_R^2) + E(\phi_L^2, \phi_R^3) - E(\phi_L^2, \phi_R^2) = 8\sqrt{2}\alpha^2\beta^2, \quad (14)$$

where $\phi_L^1 = 0$, $\phi_L^2 = \frac{\pi}{2}$, $\phi_L^3 = \frac{\pi}{4}$, and $\phi_R^1 = 0$, $\phi_R^2 = -\frac{\pi}{4}$, $\phi_R^3 = \frac{\pi}{4}$. Any local realistic theory requires $S < 2$. As Eq. (14) shows, S varies with degree of entanglement, for maximally entangled states, the quantity is $2\sqrt{2}$. For $0.479 \leq \alpha \leq 0.878$ or $-0.878 \leq \alpha \leq -0.479$, the inequality is violated.

We have a brief conclusion. In this report, we describe an experimental scheme of generating nonmaximal entanglement between two atomic ensembles. The degree of entanglement is readily tunable. This protocol fits well the status of the current experimental technology. Finally we use the nonmaximally entangled states to measure the CHSH inequality.

We thank L.-M. Duan for helpful discussion, and Y.-S. Zhang and Z.-W. Zhou for stimulating comments. This work was funded by National Fundamental Research Program (2001CB309300), National Natural Science Foundation of China, the Innovation funds from Chinese Academy of Sciences, and also by the outstanding Ph. D thesis award and the CAS’s talented scientist award entitled to Luming Duan.

-
- [1] P. W. Shor, in *Proceedings of the 35th Annual Symposium on the Foundations of Computer Science*, p. 124-133 (IEEE Computer Society Press, Los Alamitos, California, 1984).
 - [2] L. K. Grover, *Phys. Rev. Lett.* **79**, 325 (1997).
 - [3] A. K. Ekert and R. Jozsa, *Rev. Mod. Phys.* **68**, 733 (1996).
 - [4] C. H. Bennett *et al.*, *Phys. Rev. Lett.* **70**, 1895 (1993).
 - [5] D. Bouwmeester *et al.*, *Nature (London)* **390**, 575 (1997).
 - [6] C. H. Bennett and S. J. Wiesner, *Phys. Rev. Lett.* **69**, 2881 (1992).
 - [7] A. K. Ekert, *Phys. Rev. Lett.* **67**, 661 (1991).
 - [8] H.-K. Lo, H.-F. Chau, and M. Ardehali, Preprint *quant-ph/0011056*.
 - [9] P. Xue, C.-F. Li, and G.-C. Guo, *Phys. Rev. A*, **64**, 032305 (2001).
 - [10] P. Xue, C.-F. Li, and G.-C. Guo, *Phys. Rev. A*, **65**, 034302 (2002).
 - [11] P. G. Kwiat *et al.*, *Phys. Rev. Lett.* **75**, 4337 (1995).
 - [12] P. G. Kwiat, E. Waks, A. G. White, I. Appelbaum and P. H. Eberhard, *Phys. Rev. A* **60**, R773 (1999).
 - [13] A. G. White, D. F. V. James, P. H. Eberhard and P. G. Kwiat, *Phys. Rev. Lett.* **83**, 3103 (1999).
 - [14] E. Hagley *et al.*, *Phys. Rev. Lett.* **79**, 1 (1997); Q. A. Turchette *et al.*, *Phys. Rev. Lett.* **81**, 3631 (1998).
 - [15] D. Bouwmeester *et al.*, *Phys. Rev. Lett.* **82**, 1345 (1999).
 - [16] A. Rauschenbeutel *et al.*, *Science* **288**, 2024 (2000).
 - [17] C. A. Sackett *et al.*, *Nature (London)* **404**, 256 (2000).
 - [18] J. W. Pan *et al.*, *Phys. Rev. Lett.* **86**, 4435 (2001).
 - [19] M. D. Lukin *et al.*, *Phys. Rev. Lett.* **87**, 037901 (2001).
 - [20] L.-M. Duan, J. I. Cirac, P. Zoller, and E. S. Polzik, *Phys. Rev. Lett.* **85**, 5643 (2000).
 - [21] L.-M. Duan, M. D. Lukin, J. I. Cirac, and P. Zoller, *Nature (London)* **414**, 413 (2001).
 - [22] L.-M. Duan, *Phys. Rev. Lett.* **88**, 170402 (2002).
 - [23] A. Kuzmich, N. P. Bigelow, and L. Mandel, *Europhys. Lett. A* **42** 481 (1998); J. Hald, J. L. Sorensen, C. Schori, and E. S. Polzik, *Phys. Rev. Lett.* **83**, 1319 (1999); A. Kuzmich, L. Mandel, and N. P. Bigelow, *Phys. Rev. Lett.* **85**, 1594 (2000); A. Sorensen, and K. Molmer, Preprint *quant-ph/0011035*.
 - [24] B. Julsgard, A. Kozhekin, and E. S. Polzik, *Nature (London)* **413**, 400 (2001).
 - [25] P. Xue, and G.-C. Guo, Preprint *quant-ph/0205176*.
 - [26] L.-M. Duan, J. I. Cirac, and P. Zoller, Preprint *quant-*

ph/0205005.

- [27] P. H. Eberhard, Phys. Rev. Lett. **47**, R747 (1993); A. Garuccio, Phys. Rev. A **52**, 2535 (1995).
- [28] L. Hardy, Phys. Rev. Lett. **71**, 1665 (1993).
- [29] P. Xue, C.-F. Li, and G.-C. Guo, Phys. Rev. A **64**, 032304 (2001).
- [30] Q. A. Turchette *et al.*, Phys. Rev. Lett. **81**, 3631 (1998).
- [31] J. R. Torgerson, D. Branning, C. H. Monken, and L. Mandel, Phys. Lett. A **204**, 323 (1995).
- [32] G. DiGiuseppe *et al.*, Phys. Rev. A **56**, 176 (1997).
- [33] A. G. White *et al.*, Phys. Rev. Lett. **83**, 3103 (1999).
- [34] D. F. Phillips *et al.*, Phys. Rev. Lett. **86**, 783 (2001).
- [35] C. Liu, Z. Dutton, C. H. Behroozi, and L. V. Hau, Nature (London) **409**, 490 (2001).
- [36] J. F. Roch *et al.* Phys. Rev. Lett. **78**, 634 (1997).
- [37] P. Xue *et al.* unpublished.
- [38] M. Born and E. Wolf, Principles of Optics (Cambridge University press, Cambridge, 1999), 7th ed.
- [39] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23** 880 (1969).

Figure captions:

Figure 1. The relevant type Λ -level structure of the alkali atoms in the ensembles. A pair of metastable lower states $|g\rangle$ and $|s\rangle$ can be achieved, for example, Zeeman sublevels of electronic ground states $5S_{1/2}$ ^{87}Rb atoms, and $|e\rangle$ ($5P_{1/2}$) is the excited state.

Figure 2. Schematic setup for entangling two ensembles 1 and 2 in the nonmaximally entangled state. The ensembles are illuminated by the synchronized pumping laser pulses and the forward-scattering Stokes pulses are collected after the filters. The dashed line represents the pumping laser pulses with the frequency Ω and the solid line represents the Stokes pulses with the frequency ω .

Figure 3. Schematic setup for the realization of the CHSH inequality detection. Two pairs of ensembles L_1 , R_1 and L_2 , R_2 have been prepared in an EPR type of entangled state and a nonmaximally entangled state, respectively. The collective atomic excitations on each side are transferred to the optical excitations, which respectively are detected by the detectors. By choosing the relative phase shift ϕ_L and ϕ_R , we can measure the CHSH inequality.




Figure. 1




Figure. 2




Figure. 3




Figure. 2




Figure. 3